

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2003 (15.05.2003)

PCT

(10) International Publication Number
WO 03/040126 A1

(51) International Patent Classification⁷: **C07D 333/16**,
A61K 31/559, A61P 27/06

(21) International Application Number: PCT/US02/34301

(22) International Filing Date: 25 October 2002 (25.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/338,838 5 November 2001 (05.11.2001) US

(71) Applicant: **ALLERGAN, INC.** [US/US]; 2525 Dupont
Drive, Irvine, CA 92612 (US).

(72) Inventors: **BURK, Robert, M.**; 1337 Cerritos Drive,
Laguna Beach, CA 92651 (US). **HOLOBOSKI, Mark**;
29342 Bobolink Drive, Laguna Niguel, CA 92677 (US).
POSNER, Mari, F.; 27935 Hedgeline Drive, Laguna
Niguel, CA 92677 (US).

(74) Agents: **BARAN, Robert, J.** et al.; c/o Allergan, Inc.,
2525 Dupont Drive, Irvine, CA 92612 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: ω -CYCLOALKYL 17-HETEROARYL PROSTAGLANDIN E₂ ANALOGS AS EP₂-RECEPTOR AGONISTS



(I)



(II)

(57) Abstract: The invention relates to the use of derivatives of E-type prostaglandins as EP₂ agonists, in general, and, in particular as ocular hypotensives. The PGE derivatives used in accordance with the invention are represented by the formula (I): wherein the hatched segment represents an α bonds, the solid triangle represents a β bond, the wavy segments represent α or β bond, dashed lines represent a double bond or a single bond, X is selected from the group consisting of hydrogen and halogen radicals, R³ is heteroaryl or a substituted heteroaryl radical, R¹ and R² are independently selected from the group consisting of hydrogen or a lower alkyl radical having up to six carbon atoms, or a lower acyl radical having up to six carbon atoms, R is selected from the group consisting of CO₂R⁴, CONR⁴₂, CH₂OR⁴, CONR⁴SO₂R⁴, P(O)(OR⁴) and formula (II), wherein R⁴ is selected from the group consisting of H, phenyl and lower alkyl having from one to six carbon atoms and n is 0 or an integer of from 1 to 4.

WO 03/040126 A1

ω -CYCLOALKYL 17-HETEROARYL PROSTAGLANDIN
E₂ ANALOGS AS EP₂-RECEPTOR AGONISTS

5

1. Background of the Invention

2.

This application is based on, and claims the benefit of, U.S. Provisional Application No. 60/338,838, filed November 5, 2001.

10

1. Field of the Invention

The present invention relates to ω -cycloalkyl 17-heteroaryl prostaglandin E₂ analogs as EP₂-receptor agonists. These compounds are potent ocular hypotensive and are particularly suited for the management of glaucoma.

15

2. Description of Related Art

Ocular hypotensive agents are useful in the treatment of a number of various ocular hypertensive conditions, such as post-surgical and post-laser trabeculectomy ocular hypertensive episodes, glaucoma, and as presurgical adjuncts.

20

Glaucoma is a disease of the eye characterized by increased intraocular pressure. On the basis of its etiology, glaucoma has been classified as primary or secondary. For example, primary glaucoma in adults (congenital glaucoma) may be either open-angle or acute or chronic angle-closure. Secondary glaucoma results from pre-existing ocular diseases such as uveitis, intraocular tumor or an enlarged cataract.

25

The underlying causes of primary glaucoma are not yet known. The increased intraocular tension is due to the obstruction of aqueous humor outflow. In chronic open-angle glaucoma, the anterior chamber and its anatomic structures appear normal, but drainage of the aqueous humor is impeded. In acute or chronic angle-closure angle-closure glaucoma, the anterior chamber is shallow, the filtration angle is narrowed, and the iris may obstruct the trabecular meshwork at the entrance of the canal of Schlemm. Dilation of the pupil may push the root of the iris forward against the angle, and may produce pupillary block and thus

30


35

precipitate an acute attack. Eyes with narrow anterior chamber angles are predisposed to acute angle-closure glaucoma attacks of various degrees of severity.

5 Secondary glaucoma is caused by any interference with the flow of aqueous humor from the posterior chamber into the anterior chamber and subsequently, into the canal of Schlemm. Inflammatory disease of the anterior segment may prevent aqueous escape by causing complete posterior synechia in iris bombe, and may plug the drainage channel with exudates. Other common causes are intraocular tumors, enlarged cataracts, central retinal vein occlusion, 10 trauma to the eye, operative procedures and intraocular hemorrhage.

Considering all types together, glaucoma occurs in about 2% of all persons over the age of 40 and may be asymptotic for years before progressing to rapid loss of vision. In cases where surgery is not indicated, topical β -adrenoreceptor antagonists have traditionally been the drugs of choice for treating 15 glaucoma.

Certain eicosanoids and their derivatives have been reported to possess ocular hypotensive activity, and have been recommended for use in glaucoma management. Eicosanoids and derivatives include numerous biologically important compounds such as prostaglandins and their derivatives. 20 Prostaglandins can be described as derivatives of prostanoic acid which have the following structural formula:



25 Various types of prostaglandins are known, depending on the structure and substituents carried on the alicyclic ring of the prostanoic acid skeleton. Further classification is based on the number of unsaturated bonds in the side chain indicated by numerical subscripts after the generic type of prostaglandin [e.g. prostaglandin E₁ (PGE₁), prostaglandin E₂ (PGE₂)], and on the

configuration of the substituents on the alicyclic ring indicated by α or β [e.g. prostaglandin F₂ α (PGF₂ α)].

Prostaglandins were earlier regarded as potent ocular hypertensives, however, evidence accumulated in the last decade shows that some prostaglandins are highly effective ocular hypotensive agents, and are ideally suited for the long-term medical management of glaucoma (see, for example, Bito, L.Z. Biological Protection with Prostaglandins, Cohen, M.M., ed., Boca Raton, Fla, CRC Press Inc., 1985, pp. 231-252; and Bito, L.Z., Applied Pharmacology in the Medical Treatment of Glaucomas Drance, S.M. and Neufeld, A.H. eds., New York, Grune & Stratton, 1984, pp. 477-505. Such prostaglandins include PGF₂ α , PGF₁ α , PGE₂, and certain lipid-soluble esters, such as C₁ to C₂ alkyl esters, e.g. 1-isopropyl ester, of such compounds.

Although the precise mechanism is not yet known experimental results indicate that the prostaglandin-induced reduction in intraocular pressure results from increased uveoscleral outflow [Nilsson et.al., Invest. Ophthalmol. Vis. Sci. (suppl), 284 (1987)].

The isopropyl ester of PGF₂ α has been shown to have significantly greater hypotensive potency than the parent compound, presumably as a result of its more effective penetration through the cornea. In 1987, this compound was described as "the most potent ocular hypotensive agent ever reported" [see, for example, Bito, L.Z., Arch. Ophthalmol. 105, 1036 (1987), and Siebold et.al., Prodrug 5 3 (1989)].

Whereas prostaglandins appear to be devoid of significant intraocular side effects, ocular surface (conjunctival) hyperemia and foreign-body sensation have been consistently associated with the topical ocular use of such compounds, in particular PGF₂ α and its prodrugs, e.g., its 1-isopropyl ester, in humans. The clinical potentials of prostaglandins in the management of conditions associated with increased ocular pressure, e.g. glaucoma are greatly limited by these side effects.


In a series of co-pending United States patent applications assigned to Allergan, Inc. prostaglandin esters with increased ocular hypotensive activity accompanied with no or substantially reduced side-effects are disclosed. The co-pending USSN 596,430 (filed 10 October 1990), now U.S. Patent 5,446,041, relates to certain 11-acyl-prostaglandins, such as 11-pivaloyl, 11-acetyl, 11-isobutyryl, 11-valeryl, and 11-isovaleryl PGF₂ α . Intraocular pressure reducing 15-acyl prostaglandins are disclosed in the co-pending application USSN 175,476

4

(filed 29 December 1993). Similarly, 11,15- 9,15- and 9,11-diester of prostaglandins, for example 11,15-dipivaloyl PGF_{2α} are known to have ocular hypotensive activity. See the co-pending patent applications USSN Nos. 385,645 (filed 07 July 1989, now U.S. Patent 4,994,274), 584,370 (filed 18 September 1990, now U.S. Patent 5,028,624) and 585,284 (filed 18 September 1990, now U.S. Patent 5,034,413). The disclosures of all of these patent applications are hereby expressly incorporated by reference.


Summary of the Invention

The present invention concerns a method of treating ocular hypertension which comprises administering to a mammal having ocular hypertension a therapeutically effective amount of a compound of formula I



wherein the hatched segment represents an α bond, the solid triangle represents a β bond, the wavy segment represents α or β bond, dashed lines represent a double bond or a single bond, X is hydrogen or a halo radical, e.g. a fluoro or chloro radical, R³ is heteroaryl or a substituted heteroaryl radical, R¹ and R² are independently selected from the group consisting of hydrogen or a lower alkyl radical having up to six carbon atoms, or a lower acyl radical having up to six carbon atoms, R is selected from the group consisting of CO₂R⁴, CONR⁴₂, CH₂OR⁴, CONR⁴SO₂R⁴, P(O)(OR⁴) and

5



wherein R^4 is selected from the group consisting of H, phenyl and lower alkyl having from one to six carbon atoms and n is 0 or an integer of from 1 to 4.

In a further aspect, the present invention relates to an ophthalmic solution comprising a therapeutically effective amount of a compound of formula (I), wherein the symbols have the above meanings, or a pharmaceutically acceptable salt thereof, in admixture with a non-toxic, ophthalmically acceptable liquid vehicle, packaged in a container suitable for metered application. In particular, the substituents on the heteroaryl radical may be selected from the group consisting of lower alkyl, e.g. C_1 to C_6 alkyl; OR^4 ; CO_2R^4 ; halogen, e.g. fluoro, chloro and bromo; trifluoromethyl (CF_3); COR^4 , e.g. $COCH_3$; $COCF_3$; SO_2NR^4 , e.g. SO_2NH_2 ; NO_2 ; CN ; etc.

In a still further aspect, the present invention relates to a pharmaceutical product, comprising

a container adapted to dispense its contents in a metered form; and an ophthalmic solution therein, as hereinabove defined.

Finally, certain of the compounds represented by the above formula, disclosed below and utilized in the method of the present invention are novel and unobvious.

20

Brief Description of the Drawing Figures


Figure 1 is a schematic of the chemical synthesis of certain compounds related to the compounds of the invention, as specifically disclosed in Examples 12H and L and 13 H and L.

25

Figure 2 is a schematic of the chemical synthesis of certain compounds related to the compounds of the invention, as specifically disclosed in Examples 16H and L and 17H and L.

Detailed Description of the Invention

5 The present invention relates to the use of ω -cycloalkyl 17-heteroaryl
 prostaglandin E₂ analogs as EP₂-receptor agonists. The compounds used in
 accordance with the present invention are encompassed by the following
 structural formula I:




10


wherein the substituents and symbols are as hereinabove defined. The dotted lines
 on bonds between carbons 5 and 6 (C-5) and carbons 13 and 14 (C-13) indicate a
 single or double bond. If two solid lines are used at C-5, or C-13, it indicates a
 specific configuration for that double bond. Hatched lines used at position C-8,
 15 C-9 and C-11 indicate the α configuration. A triangle at position C-12 represents
 β orientation.

A preferred group of the compounds of the present invention includes
 compounds that have the following structural formula II:

20




wherein Z is selected from the group consisting of O and S, A is selected from the group consisting of N, -CH, and C, R⁵ is selected from the group consisting of hydrogen, halogen, lower alkyl having from 1 to 6 carbon atoms and lower alkoxy having from 1 to 6 carbon atoms, R⁶ and R⁷ are selected from the group consisting of hydrogen, halogen, lower alkyl having from 1 to 6 carbon atoms and lower alkoxy having from 1 to 6 carbon atoms, or, together with



, R⁶ and R⁷ forms a condensed aryl ring.

Another preferred group includes compounds having the formula III:



In the above formulae, the substituents and symbols are as hereinabove defined.

The above compounds of the present invention may be prepared by methods that are known in the art or according to the working examples below.

The compounds, below, are especially preferred representative of the compounds of the present invention.

7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-5-chloro-cyclopentyl}hept-5-enoic acid methyl ester

7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-chlorocyclopentyl}hept-5-enoic acid

(Z)-7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl}hept-5-enoic acid methyl ester

(Z)-7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl}hept-5-enoic acid

5 (Z)-7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]cyclopentenyl}hept-5-enoic acid methyl ester

10 (Z)-7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]cyclopentenyl}hept-5-enoic acid

15 A pharmaceutically acceptable salt is any salt which retains the activity of the parent compound and does not impart any deleterious or undesirable effect on the subject to whom it is administered and in the context in which it is administered. Of particular interest are salts formed with inorganic ions, such as sodium, potassium, calcium, magnesium and zinc.

20 Pharmaceutical compositions may be prepared by combining a therapeutically effective amount of at least one compound according to the present invention, or a pharmaceutically acceptable acid addition salt thereof, as an active ingredient, with conventional ophthalmically acceptable pharmaceutical excipients, and by preparation of unit dosage forms suitable for topical ocular use. The therapeutically efficient amount typically is between about 0.0001 and about 5% (w/v), preferably about 0.001 to about 1.0% (w/v) in liquid formulations.

25 For ophthalmic application, preferably solutions are prepared using a physiological saline solution as a major vehicle. The pH of such ophthalmic solutions should preferably be maintained between 6.5 and 7.2 with an appropriate buffer system. The formulations may also contain conventional, pharmaceutically acceptable preservatives, stabilizers and surfactants.

30 Preferred preservatives that may be used in the pharmaceutical compositions of the present invention include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate. A preferred surfactant is, for example, Tween 80. Likewise, various preferred vehicles may be used in the ophthalmic preparations of the present invention. These vehicles include, but are not limited to, polyvinyl alcohol, povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose and purified water.

35 Tonicity adjustors may be added as needed or convenient. They include, but are not limited to, salts, particularly sodium chloride, potassium chloride,

mannitol and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

In a similar vein, an ophthalmically acceptable antioxidant for use in the present invention includes, but is not limited to, sodium metabisulfite, sodium thiosulfate, acetylcysteine, butylated hydroxyanisole and butylated hydroxytoluene.

Other excipient components which may be included in the ophthalmic preparations are chelating agents. The preferred chelating agent is edentate disodium, although other chelating agents may also be used in place or in conjunction with it.

The ingredients are usually used in the following amounts:

	<u>Ingredient</u>	<u>Amount (% w/v)</u>
	active ingredient	about 0.001-5
	preservative	0-0.10
	vehicle	0-40
	tonicity adjustor	1-10
	buffer	0.01-10
	pH adjustor	q.s. pH 4.5-7.5
	antioxidant	as needed
	surfactant	as needed
	purified water	as needed to make 100%

The actual dose of the active compounds of the present invention depends on the specific compound, and on the condition to be treated; the selection of the appropriate dose is well within the knowledge of the skilled artisan.

The ophthalmic formulations of the present invention are conveniently packaged in forms suitable for metered application, such as in containers equipped with a dropper, to facilitate the application to the eye. Containers suitable for dropwise application are usually made of suitable inert, non-toxic plastic material, and generally contain between about 0.5 and about 15 ml solution.

The invention is further illustrated by the following non-limiting Examples, which are summarized in the reaction schemes of Figures 1 and 2,

10

wherein the compounds are identified by the same designator in both the Examples and the Figures.

Example 1

Ethyl cyclobutanecarboxylate acid ethyl ester (1).

The named compound was purchased from Aldrich Chemical Co., P.O. Box 2060, Milwaukee, WI 53201 USA.

Example 2

1-(1-Hydroxy-1-thiophen-2-yl-methyl)cyclobutanecarboxylic acid ethyl ester (2).

Lithium diisopropylamide mono(THF) (1.95 mL of a 2.0M solution in heptane/THF/ethylbenzene, 3.90 mmol) was added to a solution of ester 1 (0.50 g, 3.9 mmol) in THF (6 mL) at -78°C . After stirring 30 min, 2-thiophenecarboxaldehyde (667 mg, 5.95 mmol) was added and the mixture was stirred for 3 h. After the reaction was judged complete by TLC analysis, saturated aqueous NH_4Cl was added and the reaction was slowly warmed to 23°C . The THF was evaporated and the reaction mixture was extracted with CH_2Cl_2 (2x). The combined organic layers were washed with brine, dried (Na_2SO_4), filtered and concentrated *in vacuo*. Purification of the residue by flash column chromatography (FCC) (silica gel, 100% hexane followed by 9 : 1 hexane / EtOAc) afforded the above named compound 2.

Example 3

1-Thiophen-2-yl-methylcyclobutanecarboxylic acid ethyl ester (3).

Trimethylsilyliodide (20 g, 100 mmol) was added to CH_3CN (10 mL) at 0°C and the mixture was allowed to stir 5 min. A solution of alcohol 2 (5 g, 20 mmol) in CH_3CN (10 mL) was added slowly while the temperature was kept between $4-10^{\circ}\text{C}$ and the reaction was allowed to warm to 23°C . After stirring for 2 h at 23°C the reaction was judged complete *via* TLC analysis. The mixture was poured into 3N NaOH at 0°C and EtOAc was added. The organic layer was separated, washed with brine, dried (Na_2SO_4), filtered and

11

concentrated *in vacuo*. FCC (silica gel, 1 : 1 hexane / CH₂Cl₂) gave 2.3 g of the above named ester 3.

Example 4

5

(1-Thiophen-2-yl-methylcyclobutyl)methanol (4).

Lithium borohydride (435 mg, 20 mmol) was added to a solution of ester 3 (2.3 g, 10 mmol) in Et₂O (20 mL) at 0 °C. After having stirred for 5 min, MeOH (640 mg, 20 mmol) was added dropwise and stirring continued at 0 °C until effervescence ceased. The mixture was warmed to 23 °C and was allowed to stir an additional hour, at which time the mixture was poured into 3N NaOH and stirred an additional 0.5 h. The organic layer was separated and washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. The crude alcohol 4 was purified by FCC (silica gel, 1 : 1 hexane / CH₂Cl₂).

20

Example 5

1-Thiophen-2-yl-methylcyclobutanecarbaldehyde (5).

Oxalyl chloride (50 mL, 0.10 mmol) was added to CH₂Cl₂ (150 mL) at 23 °C and was cooled to -78 °C. DMSO (16 g, 0.20 mmol) was added dropwise to the mixture and stirring was continued for 15 min. A solution of alcohol 4 (7.9 g, 0.041 mmol) in CH₂Cl₂ (50 mL) was then added dropwise, after which Et₃N (44 g, 0.44 mmol) was added and the mixture was warmed to 23 °C. After 1 h, the mixture was poured into saturated aqueous NaHCO₃ and the organic layer was separated. The aqueous layer was extracted with CH₂Cl₂ (2x) and the combined organic portions were washed with brine, dried (Na₂SO₄), concentrated *in vacuo* and purified by FCC (silica gel, 100% hexane followed by 2 : 1 hexane / CH₂Cl₂) to afford the above named aldehyde 5.

Example 6**1-(1-Thiophen-2-yl-methylcyclobutyl)but-2-yn-1-ol (6).**

5 A solution of propylmagnesium bromide (360 mL of a 0.5M solution in THF, 0.180 mmol; 0.5 M in THF) was added dropwise to a solution of aldehyde **5** (7.0 g, 36 mmol) in THF (200 mL) while the mixture was maintained at ambient temperature. After having stirred 3 h at 23 °C, the reaction was poured into saturated aqueous NH₄Cl and extracted with Et₂O. The organic portion was
10 separated and was washed with saturated aqueous NaHCO₃, brine, then dried (Na₂SO₄) and concentrated *in vacuo*. FCC (silica gel, 100% hexane followed by 1 : 1, hexane / CH₂Cl₂) gave 6.2 g of the above named alkyne **6**.

Example 7

15

1-(1-Thiophen-2-yl-methylcyclobutyl)but-3-yn-1-ol (7).

A dry round bottom flask was charged with potassium hydride (5.5 g, 48 mmol; 35% by wt dispersion in oil) and the oil was removed by hexane rinse (3x).
20 Aminopropylamide (39 mL) was added to the mixture and it was stirred until effervescence ceased. The mixture was then cooled to 0 °C and the alkyne **6** (2 g, 9.1 mmol) was added and the reaction stirred at 23 °C for 1 h. The reaction was quenched with MeOH (2 mL) and water. The mixture was extracted with Et₂O (3x) and the combined organic layer was washed with 1N HCl, brine, dried
25 (Na₂SO₄) and concentrated *in vacuo*. FCC (silica gel, 1 : 1 hexane / CH₂Cl₂) gave 570 mg of the above named alkyne **7**.

Example 8

30 **tert-Butyldimethyl[1-(1-thiophen-2-yl-methylcyclobutyl)but-3-ynyloxy]silane (8).**

13

To a cooled (0 °C) solution of alkyne **7** (200 mg, 0.9 mmol), CH₂Cl₂ (5 mL) and triethylamine (275 mg, 2.72 mmol) was added *tert*-butyldimethylsilyl trifluoromethanesulfonate (360 mg, 1.36 mmol) dropwise. After having stirred for 5 min at 0 °C, the mixture was warmed to 23 °C and stirred an additional hour. The reaction was then quenched with saturated aqueous NaHCO₃ and extracted with CH₂Cl₂ (2X). The combined organics were washed with 1N HCl, saturated aqueous NaHCO₃, brine then were dried (Na₂SO₄), filtered and concentrated *in vacuo*. FCC (silica gel, 100% hexane) gave 695 mg of the above named compound **8**.

Example 9

***tert*-Butyl-[(E)-4-iodo-1-(1-thiophen-2-ylmethylcyclobutyl)but-3-enyloxy]dimethylsilane (9).**

Cp₂ZrHCl (304 mg, 1.18 mmol) was added to a solution of alkyne **8** (263 mg, 0.786 mmol) in CH₂Cl₂ (5 mL) at 23 °C and stirring was maintained for 20 min. N-iodosuccinimide (247 mg, 1.18 mmol) was added to the mixture and stirring was continued for an additional 30 min. The mixture was concentrated *in vacuo*, diluted with hexane / Et₂O, filtered and concentrated *in vacuo*. FCC (silica gel, 100% hexane) gave 360 mg of the above named compound **9**.

Example 10

7-[(R)-3-(*tert*-Butyldimethylsilyloxy)-5-oxo-cyclopent-1-enyl]heptanoic acid methyl ester (10).

The named compound was purchased from Nissan Chemical Industries, LTD, Tokyo 101-0054 Japan.

Example 11

7-[(1R,2R,3R)-2-[(E)-(*tert*-Butyldimethylsilyloxy)-(1-thiophen-2-ylmethylcyclobutyl)but-1-enyl]-3-[(dimethylethyl)dimethylsilyloxy]-5-oxo-cyclopentyl]heptanoic acid methyl ester (11).

To a solution of vinyl iodide **9** (120 mg, 0.259 mmol) in Et₂O (1.5 mL) at

14

–78 °C was added *t*-BuLi (0.35 mL of a 1.5M solution in THF, 0.52 mmol). After the mixture had stirred 30 min at –78 °C 2-thienyl(cyano)copper lithium (1.14 mL, 0.285 mmol) was added and stirring was continued for an additional 30 min. The reaction was then treated with a solution of the enone 10 (91.6 mg, 0.259 mmol) in Et₂O (1 mL). After several minutes had passed, the reaction had solidified and 0.5 mL Et₂O was added. The reaction was stirred 1 h at –78 °C, was poured into saturated aqueous NH₄Cl and then was extracted with EtOAc (3x). The combined organic portions were washed with brine, filtered and concentrated *in vacuo*. FCC (silica gel, 100% hexane; 9 : 1 hexane / EtOAc) gave 63 mg of the above named compound 11.

Example 12

7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-thiophen-2-yl-methyl-cyclobutyl)but-1-enyl]-5-oxo-cyclopentyl}heptanoic acid methyl ester (12H).

7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-thiophen-2-yl-methyl-cyclobutyl)but-1-enyl]-5-oxo-cyclopentyl}heptanoic acid methyl ester (12L).

Hydrogen fluoride-pyridine (0.091 mL) was added to a solution of the *bis*-TBS ether 11 (63 mg, 0.912 mmol) in CH₃CN (3 mL) at 23 °C. After having stirred for 3 h, the mixture was quenched with saturated aqueous Na₂CO₃ and extracted with EtOAc (3x). The combined organic portions were washed with 1N HCl, saturated aqueous NaHCO₃, brine, and were then dried (Na₂SO₄), filtered and concentrated *in vacuo*. FCC (silica gel, 3 : 2 hexane / EtOAc followed by 1 : 1 hexane / EtOAc) gave a higher R_f diol (10 mg) and a lower R_f diol (30 mg), hereinafter, designated as named compounds 12H and 12L, respectively).

Example 13H

7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-thiophen-2-yl-methylcyclobutyl)but-1-enyl]-5-oxocyclopentyl}heptanoic acid (13H).

15

Methyl ester **12H** (4.8 mg, 10.4 mmol) and PLE (0.134 mmol, 45 mmol) were stirred in phosphate buffer (2 mL, pH 7.2) at 23 °C over 16 h. After the reaction was complete, the mixture was filtered and the aqueous phase was extracted with EtOAc (3x). The combined organic phases were washed with brine, dried
5 (Na₂SO₄), filtered and concentrated *in vacuo*. FCC (silica gel, 1 : 1 hexane / EtOAc; EtOAc) gave of the above named acid **13H**.

Example 13L

10 **7-[(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-thiophen-2-yl-methyl-cyclobutyl)but-1-enyl]-5-oxo-cyclopentyl]heptanoic acid (13L).**

Methyl ester **12L** was reacted according to Example 13H to yield the above named compound.

15

Example 14

(Z)-7-[(R)-3-(tert-Butyldimethylsilanyloxy)-5-oxo-cyclopent-1-enyl]hept-5-enoic acid methyl ester (14).

20

The named compound was purchased from Nissan Chemical Industries, LTD, Tokyo 101-0054 Japan.

Example 15

25

(Z)-7-[(1R,2R,3R)-2-[(E)-(tert-Butyldimethylsilanyloxy)-(1-thiophen-2-yl-methylcyclobutyl)but-1-enyl]-3-[(dimethylethyl)dimethylsilanyloxy]-5-oxo-cyclopentyl]hept-5-enoic acid methyl ester (15).

30

The compound of Example 14, above, was reacted in accordance with the process of Example 11 to yield the above named compound.

Example 16

16

(Z)-7-[(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-thiophen-2-yl-methylcyclobutyl)but-1-enyl]-5-oxo-cyclopentyl]hept-5-enoic acid methyl ester (16H).

5 (Z)-7-[(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-thiophen-2-yl-methylcyclobutyl)but-1-enyl]-5-oxo-cyclopentyl]hept-5-enoic acid methyl ester (16L).

10 The compound of Example 15 is reacted in accordance with the process of Example 12 to yield a higher R_f diol (6.0 mg) and a lower R_f diol (6.0 mg), hereinafter, designated as 16H and 16L, respectively.

15 Example 17H

(Z)-7-[(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-thiophen-2-yl-methylcyclobutyl)but-1-enyl]-5-oxo-cyclopentyl]hept-5-enoic acid (17H).

20 The compound 16H of Example 16 is reacted in accordance with the process of Example 13H to yield the above named compound.

Example 17L

25 (Z)-7-[(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-thiophen-2-yl-methylcyclobutyl)but-1-enyl]-5-oxo-cyclopentyl]hept-5-enoic acid (17L).

30 The compound 16L of Example 16 is reacted in accordance with the process of Example 13H to yield the above named compound.

Example 18

17

7-**{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-5-chloro-cyclopentyl}**hept-5-enoic acid methyl ester (18H and 18L).

- 5 Examples 14 through 16 is repeated with the appropriate chloro derivative replacing (Z)-7-[(R)-3-(*tert*-Butyldimethylsilanyloxy)-5-oxo-cyclopent-1-enyl]hept-5-enoic acid methyl ester **14** and the appropriate chlorothieryl derivative replacing 2-thienyl(cyano)copper lithium to yield a product which is separated to provide a higher R_f diol and a lower R_f diol designated as 18H and
10 18L, respectively.

Example 19H

- 15 7-**{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-5-chloro-cyclopentyl}**hept-5-enoic acid (**19H**).

- 20 The compound of Example 18H is reacted according to the process of Example 13H to yield the named compound.

Example 19L

- 25 7-**{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-5-chloro-cyclopentyl}**hept-5-enoic acid (**19L**).

- 30 The compound of Example 18L is reacted according to the process of Example 13L to yield the named compound.

Example 20

18

7-**{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl}hept-5-enoic acid methyl ester (20H and 20L).**

5 Examples 14 through 16 is repeated with the appropriate fluoro derivative replacing (Z)-7-[(R)-3-(*tert*-Butyldimethylsilanyloxy)-5-oxo-cyclopent-1-enyl]hept-5-enoic acid methyl ester 14 and the appropriate chlorothieryl derivative replacing 2-thienyl(cyano)copper lithium to yield a product which is separated to provide a higher R_f diol and a lower R_f diol designated as 20H and
10 20L, respectively.

Example 21H

15 7-**{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl}hept-5-enoic acid (21H).**

The compound of Example 20H is reacted according to the process of Example 13H to yield the named compound.

20

Example 21L

25 7-**{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl}hept-5-enoic acid (21L).**

25

The compound of Example 20L is reacted according to the process of Example 13L to yield the named compound.

Example 22

30

19

7-{{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-cyclopentenyl}hept-5-enoic acid methyl ester.

5 Examples 14 through 16 is repeated with the appropriate nor keto derivative replacing (Z)-7-[(R)-3-(*tert*-Butyldimethylsilanyloxy)-5-oxo-cyclopent-1-enyl]hept-5-enoic acid methyl ester **14** and the appropriate chlorothieryl derivative replacing 2-thienyl(cyano)copper lithium to yield a product which is separated to provide a higher R_f diol and a lower R_f diol designated as 22H and
10 22L, respectively.

Example 23H

15 7-{{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-cyclopentenyl}hept-5-enoic acid (23H).

The compound of Example 22H is reacted according to the process of Example 13H to yield the named compound.

20 Example 23L

7-{{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methyl-cyclobutyl)but-1-enyl]-cyclopentenyl}hept-5-enoic acid (23L).

25 The compound of Example 22L is reacted according to the process of Example 13L to yield the named compound.

RADIOLIGAND BINDING






30 Recombinant EP₂ receptor; transient transfectants COS-7 cells were transiently transfected using Lipofectin (Gibco-BRL life Technologies, Gaitherburg, MD, U.S.A.) according to manufacturer's protocols. For binding studies, 2×10^6 cells were plated onto 150 mm dishes 24 h prior to transfection.

20










Each plate was transfected with 50 µg plasmid DNA and 50 µL lipofectin. Cells were collected and membranes prepared at 48 h post-transfection, and frozen at -80 °C until use.

Plasma membrane preparations were thawed at room temperature and used at a final 1 mg/mL concentration in a 500 µL volume. The binding of [3H]-PGE₂ (specific activity 180 Ci mmol⁻¹) were determined in duplicate and experiments were replicated three times. Incubations were for 60 min at 25°C and were terminated by the addition of 4 mL of ice-cold 50 µM TRIS-HCl, followed by rapid filtration through Whatman GF/B filters and three additional 4 mL washes in a cell harvester (Brandel). Competition studies were performed using a final concentration of 5 nM [3H]-PGE₂ and non-specific binding determined with 10 µM of the respective unlabelled prostanoid.

Certain of the above compounds were tested for activity in the recombinant human EP₂ receptor assay described above and the results are reported in Table 1, below. Note Examples 20 and 21 are the unseparated mixtures of Examples 20H and 20L and 21H and 21L, respectively.

Example #	Structure	hEP ₂
18H		4300
18L		5500
19H		126
19L		300
22H		NA

21

22L		NA
23H		700
23L		2500
20H		5100
20L		NA
21H		132
21L		300
20		8100
21		112

EP₂ activity indicates that the compounds of this invention are useful in treating asthma, dysmenorrhea as well as glaucoma and lowering intraocular pressure.

5 Other potential therapeutic applications are in osteoporosis, constipation, renal disorders, sexual dysfunction, baldness, diabetes, cancer and in disorder of immune regulation.

The compounds of the invention may also be useful in the treatment of various pathophysiological diseases including acute myocardial infarction, vascular thrombosis, hypertension, pulmonary hypertension, ischemic heart
10 disease, congestive heart failure, and angina pectoris, in which case the compounds may be administered by any means that effect vasodilation and

thereby relieve the symptoms of the disease. For example, administration may be by oral, transdermal, parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, or buccal routes.

5 The compounds of the invention may be formulated into an ointment containing about 0.10 to 10% of the active ingredient in a suitable base of, for example, white petrolatum, mineral oil and petroatum and lanolin alcohol. Other suitable bases will be readily apparent to those skilled in the art.

10 The pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, for example, by means of conventional dissolving or suspending the compounds, which are all either water soluble or suspendable. For administration in the treatment of the other mentioned pathophysiological disorders. The pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules make of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules can contain the active compounds in liquid form
15 that may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as in buffered salt solution. In addition, stabilizers may be added.
20

In addition to being provided in a liquid form, for example in gelatin capsule or other suitable vehicle, the pharmaceutical preparations may contain suitable excipients to facilitate the processing of the active compounds into preparations that can be used pharmaceutically. Thus, pharmaceutical
25 preparations for oral use can be obtained by adhering the solution of the active compounds to a solid support, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

Suitable excipients are, in particular, fillers such as sugars, for example
30 lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, as well as inders such as starch, paste using for example, maize starch, wheat starch, rich starchy, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or
35 polyvinyl pyrrolidone. If desired, disintegrating agents may be added such as the above-mentioned starches and also carboxymethyl-starch, crosslinked


polyvinyl pyrrolidone, agar, or algenic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings which if desired, are resistant to gastric juices. For this purpose, concentrated sugar solutions may be used, which may optionally containing gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations such as acetylcellulose phthalate or hydroxypropylmethyl-cellulose phthalate, are used. Dye stuffs or pigments may be added to the tables or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

Suitable formulations for intravenous or parenteral administration include aqueous solutions of the active compounds. In addition, suspensions of the active compounds as oily injection suspensions may be administered. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, soribitol, and/or dextran. Optionally, the suspension may also contain stabilizers.


The foregoing description details specific methods and compositions that can be employed to practice the present invention, and represents the best mode contemplated. However, it is apparent for one of ordinary skill in the art that further compounds with the desired pharmacological properties can be prepared in an analogous manner, and that the disclosed compounds can also be obtained from different starting compounds via different chemical reactions. Similarly, different pharmaceutical compositions may be prepared and used with substantially the same result. Thus, however detailed the foregoing may appear in text, it should not be construed as limiting the overall scope hereof; rather, the ambit of the present invention is to be governed only by the lawful construction of the appended claims.

CLAIMS

- 5 1. A method of treating ocular hypertension which comprises administering to a mammal having ocular hypertension a therapeutically effective amount of a compound represented by formula I:



- 10 wherein the hatched segment represents an α bond, the solid triangle represents a β bond, the wavy segment represents α or β bond, dashed lines represent a double bond or a single bond, X is selected from the group consisting of hydrogen and halogen radicals, R^3 is heteroaryl or a substituted heteroaryl radical, R^1 and R^2 are independently selected from the group consisting of hydrogen or a lower alkyl radical having up to six carbon atoms, or a lower acyl radical having up to six
 15 carbon atoms, R is selected from the group consisting of CO_2R^4 , $CONR^4_2$, CH_2OR^4 , $CONR^4SO_2R^4$, $P(O)(OR^4)$ and




- wherein R^4 is selected from the group consisting of H, phenyl and lower alkyl
 20 having from one to six carbon atoms and n is 0 or an integer of from 1 to 4.

25


2. The method of Claim 1 wherein the substituent on the heteroaryl radical is selected from the group consisting of C₁ to C₆ alkyl; OR⁴; CO₂R⁴; halogen, trifluoromethyl; COR⁴; COCF₃; SO₂NR⁴; NO₂; and CN.

5 3. The method of Claim 2 wherein said compound is represented by formula II:



10 wherein Z is selected from the group consisting of O and S, A is selected from the group consisting of N, -CH, and C, R⁵ is selected from the group consisting of hydrogen, halogen, lower alkyl having from 1 to 6 carbon atoms, and lower alkoxy having from 1 to 6 carbon atoms, R⁶ and R⁷ are selected from the group consisting of hydrogen, halogen, lower alkyl having from 1 to 6 carbon atoms, lower alkoxy having from 1 to 6 carbon atoms or, together with


15



, R⁶ and R⁷ forms a condensed aryl ring.

20 4. The method of claim 3 wherein said compound is represented by formula III:

26




5. The method of claim 4 wherein the dashed line represents a double bond.
6. The method of claim 4 wherein the dashed line represents a single bond.
- 5 7. The method of claim 5 wherein R is CO₂R⁴ and R⁴ is H.
8. The method of claim 6 wherein R is CO₂R⁴ and R⁴ is H.
- 10 9. The method of claim 5 wherein R is CO₂R⁴ and R⁴ is methyl.
10. The method of claim 6 wherein R is CO₂R⁴ and R⁴ is methyl.
11. The method of claim 4 wherein R is CO₂R⁴.
- 15 12. The method of claim 11 wherein R⁴ is selected from the group consisting of H and lower alkyl.
13. The method of claim 4 wherein A is C and R₅, R₆ and R₇ are H.
- 20 14. The method of claim 1 wherein X is selected from the group consisting of hydrogen, fluoro and chloro radicals.
15. The method of claim 14 wherein X is hydrogen.
- 25 16. The method of claim 14 wherein X is fluoro.
17. The method of claim 14 wherein X is chloro.


27

18. The method of claim 1 wherein said compound is 7-{{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)methyl-cyclobutyl)but-1-enyl]-5-chloro-cyclopentyl}hept-5-enoic acid methyl ester.
- 5 19. The method of claim 1 wherein said compound is 7-{{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)methylcyclobutyl)but-1-enyl]-5-chlorocyclopentyl}hept-5-enoic acid.
- 10 20. The method of claim 1 wherein said compound is (Z)-7-{{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl}hept-5-enoic acid methyl ester
- 15 21. The method of claim 1 wherein said compound is (Z)-7-{{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl}hept-5-enoic acid.
- 20 22. An ophthalmic solution comprising a therapeutically effective amount of a compound of formula I, as defined in Claim 1, or a pharmaceutically acceptable salt thereof, in admixture with a non-toxic, ophthalmically acceptable liquid vehicle, packaged in a container suitable for metered application.
- 25 23. The ophthalmic solution of Claim 22 wherein said compound is a compound of Formula III.
- 30 24. A pharmaceutical product, comprising a container adapted to dispense the contents of said container in metered form; and an ophthalmic solution in said container comprising a compound of formula I as defined in Claim 1, or a pharmaceutically acceptable salt thereof, in admixture with a non-toxic, ophthalmically acceptable liquid vehicle.
25. The product of claim 24 wherein said compound is a compound of Formula III.
- 35 26. The compound represented by formula I:

28



- wherein the hatched segment represents an α bond, the solid triangle represents a β bond, the wavy segment represents an α or β bond, dashed lines represent a double bond or a single bond, X is selected from the group consisting of hydrogen and halogen radicals, R^3 is heteroaryl or a substituted heteroaryl radical, R^1 and R^2 are independently selected from the group consisting of hydrogen or a lower alkyl radical having up to six carbon atoms, or a lower acyl radical having up to six carbon atoms, R is selected from the group consisting of CO_2R^4 , $CONR^4_2$, CH_2OR^4 , $CONR^4SO_2R^4$, $P(O)(OR^4)$ and



- wherein R^4 is selected from the group consisting of H, phenyl and lower alkyl having from one to six carbon atoms and n is 0 or an integer of from 1 to 4.

15


27. The compound of claim 26 wherein R is CO_2R^4 .

28. The compound of claim 27 wherein R^4 is selected from the group consisting of H and lower alkyl.

20

29. The compound of claim 28 wherein said compound is represented by the formula

29



30. The compound of claim 29 wherein A is C and R₅, R₆ and R₇ are H.

31. The compound of claim 30 wherein said compound is selected from the group consisting of :

7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-chloro-cyclopentyl} hept-5-enoic acid methyl ester,


7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-chlorocyclopentyl} hept-5-enoic acid,

(Z)-7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl} hept-5-enoic acid methyl ester,

(Z)-7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]-5-fluoro-cyclopentyl} hept-5-enoic acid,

(Z)-7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]cyclopentenyl} hept-5-enoic acid methyl ester and

(Z)-7-{(1R,2R,3R)-3-Hydroxy-2-[(E)-4-hydroxy-4-(1-(5-chloro-thiophen-2-yl)-methylcyclobutyl)but-1-enyl]cyclopentenyl} hept-5-enoic acid.

1/2
FIG. 1.

2/2

FIG. 2.